

## DISINFECTANT CLEANER QUICK • EFFECTIVE • CONVENIENT CLEANS, DISINFECTS & DEODORIZES

- Hospital disinfectant ready-to-use spray
- · Disinfects in just 2 minutes.
- Kills Rhinovirus, Norovirus, Canine Parvovirus in 2 minutes!
- Kills 99.9% of Staphylococcus aureus (Staph), Klebsiella pneumoniae in just 5 seconds!
- Performex RTU has demonstrated effectiveness against Influenza A virus and is expected to inactivate all Influenza A viruses including Pandemic 2009 H1N1 Influenza A virus.
- · Use on hard, non-porous surfaces
- Do not use on unpainted wood

Performex RTU is bactericidal in the presence of organic soil (5% blood serum) against Salmonella enterica, Pseudomonas aeruginosa, Staphylococcus aureus (Staph), Methicillin Resistant Staphylococcus aureus (MRSA) and Vancomycin Resistant Enterococcus faecium (VRE) at 2 minutes contact time.

Performex RTU is intended for use in:

· Hospitals and healthcare settings: ambulances, Dental Offices, Nursing homes, Operating rooms, Physical therapy, Respiratory therapy

· Veterinary clinics, animal hospitals, kennels, zoos

- · Beauty salons, barber shops, correctional facilities, mortuaries
- Schools, gyms, athletic facilities
- Grocery stores
  Institutional kitchens and/or bathrooms

Use Performex RTU to clean and disinfect non-porous personal protective safety equipment, half mask respirators, full face breathing apparatus, gas masks, face shields, silicone rubber or PVC hearing protectors and vinyl covered ear muffs. Rinse all equipment that comes in prolonged contact with skin before reuse with clean warm water (about 120°F), and allow to air dry. (Caution: Cleaning at the 120°F temperature will avoid overheating and distortion of the personal safety equipment that would necessitate replacement.)

Performex RTU is a one-step hospital-use germicidal cleaner and deodorant designed for general cleaning, and disinfecting hard, non-porous inanimate surfaces. Quickly removes dirt, grime, food residue, blood and other organic matter commonly found in health care facilities. It also eliminates odors leaving surfaces smelling clean and fresh. Use where odors are a problem.

- Animal Housing Facilities and Poultry and Swine Premises: Performex RTU
  removes dirt, grime, blood, urine, fecal matter and other common soils found
  in animal housing facilities, grooming facilities, kennels, pet stores, veterinary clinics, laboratories or other small animal facilities. It eliminates odors leaving surfaces smelling clean and fresh.
- Non-Acid Bowl & Bathroom Disinfectant Cleaner: Performex RTU is also a ready-to-use non-acid bowl and bathroom cleaner that cleans, disinfects and deodorizes in one easy step. It cleans, disinfects and deodorizes toilet bowls, urinals, rims, sinks, sink basins, faucets, tubs, glazed tiles, glazed ceramic, glazed porcelain, chrome, stainless steel, and all non-porous inanimate surfaces found in the restroom.

### DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

This product is not to be used as a terminal sterilant/high-level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the bloodstream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body.

DISINFECTING, DEODORIZING: Remove gross filth or heavy soil prior to application of Performex RTU. Apply product with a hand pump trigger sprayer. Hold sprayer six to eight inches from the surface to be treated. Spray area until it is covered with the product. Allow surface to remain visibly wet for 2 minutes. No scrubbing necessary. Allow treated surfaces to air dry or wipe off with a clean cloth.

When using on food contact surfaces: counter tops, kitchen appliances, thoroughly rinse all treated surfaces with potable water. This product must not be used to clean eating utensils, glassware and dishes.

Performex RTU is an effective disinfectant against: Pseudomonas aeruginosa, Salmonella enterica, Staphylococcus aureus Methicillin Resistant Staphylococcus aureus (MRSA), Vancomycin Resistant Enterococcus faecium, Multi-Drug Resistant (MDR) Acinetobacter baumannii Ceftazidime and Gentimicin Resistant, Extended Spectrum Beta-Lactamase (ESBL) Escherichia coli and Klebsiella pneumoniae New Dehli Metallo-Beta Lactamase (NDM-1) Carbapenem

One-step cleaning and disinfecting of toilet bowls and urinals: Remove water from bowl by forcing water over the trap with swab mop applicator. Press swab mop against side of the bowl to remove excess water from applicator. Apply 1 to 2 ounces of this product evenly across bowl surface, allowing sufficient time for Performex RTU to cover entire surface for maximum cleaning. Swab bowl completely using a scrub brush or toilet mop, making sure to get under the rim. Let stand for 2 minutes and flush.

To disinfect food service establishment food contact surfaces: countertops, appliances, tables, hold container six to eight inches from the surface to be treated and spray area until it is covered with the solution. Allow Performex RTU to penetrate and the surface to remain wet for 2 minutes. Wipe off with a clean cloth, mop or sponge and thoroughly rinse all food contact surfaces in the area with potable water. Heavily soiled areas must be pre-cleaned before using this product. Do not use on eating utensils, glassware and cookware.

Disinfection of animal guarters and kennels: To disinfect pre-cleaned animal quarters and kennels, apply Performex RTU. Remove all animals and feed from premises. Remove all litter and droppings from floors, walls and surfaces of facilities occupied or traversed by animals. Empty all troughs, racks, and other feeding and watering appliances. Thoroughly clean all surfaces with soap or detergent and rinse with water. Saturate the surfaces with the disinfecting solution for a period of 2 minutes (for Canine Parvovirus). Ventilate building and other closed spaces. Do not house animals or employ equipment until treatment has been absorbed, set or dried. All treated equipment that will contact feed or drinking water must be rinsed with potable water before reuse.

\*VIRUCIDAL ACTIVITY: Performex RTU is an effective virucide on hard nonporous surfaces against Human Immunodeficiency Virus Type-1 (HIV-1) and Avian Influenza Virus (H5N1) NIBRG-14, when the treated surface is allowed to remain wet for 30 seconds. Performex RTU is an effective virucide on hard, non-porous surfaces against Human Rotavirus, Hepatitis B Virus (HBV), Hepatitis C Virus (HCV), Duck Hepatitis B Virus (DHBV) and Bovine Viral Diarrhea Virus (BVDV) when the treated surface is allowed to remain wet for (1 minute) (60 seconds). Performex RTU is an effective virucide on hard, non-porous surfaces against Feline Calicivirus, Norovirus (Norwalk Virus), Canine Parvovirus, Rhinovirus Type 14 and Rhinovirus Type 39 when the treated surface is allowed to remain wet for 2 minutes.

KILLS HIV-1, HBV and HCV ON PRE-CLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS in health care settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/ objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1, Hepatitis B Virus (HBV) and Hepatitis C Virus (HCV).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1, HBV AND HCV OF SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS.

PERSONAL PROTECTION: When handling items soiled with blood or body fluids, use disposable latex gloves, gowns, masks, and eye coverings.

CLEANING PROCEDURES: Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of Performex RTU.

CONTACT TIME: Allow surfaces to remain wet for 2 minutes. (HIV-1 is inactivated in 30 seconds, HBV and HCV are inactivated in 1 minute).

DISPOSAL OF INFECTIOUS MATERIALS: Blood and other body fluids must be autoclaved and disposed of according to local regulations for infectious waste disposal.

SANITIZE NON-FOOD CONTACT SURFACES and DEODORIZE: To sanitize/ deodorize hard, non-porous surfaces, hold container six to eight inches from the surface to be treated. Spray area until it is covered with the product. Allow surface to remain visibly wet for 5 seconds. Performex RTU is an effective sanitizer against Staphylococcus aureus (Staph) and Klebsiella pneumoniae. Let surface dry. For heavily soiled surfaces, a pre-cleaning step is required.

For surfaces that may come in contact with food, a potable water rinse is required.

# PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

## FIRST AID

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

IF IN EYES: Hold eyes open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice. IF ON SKIN: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment

### STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in a dry place inaccessible to children.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available or puncture and dispose in a sanitary landfill, incineration, or burning, if allowed by state and local authorities. If burned,

stay out of smoke.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

EPA Reg. No. 1839-220-106 **EPA 11 MAY 15** 

EPA Est. No. 106-IN-1 LN:4032